arXiv:astro-ph/0212363v2 16 Apr 2003

The Amplitude of Mass Fluctuations

Neta A. Bahcall and Paul Bode

Princeton University Observatory, Princeton, NJ 08544-1001

neta@astro.princeton.edu, bode@astro.princeton.edu

ABSTRACT

We determine the linear amplitude of mass fluctuations in the universe, σ_8, from the abundance of massive clusters at redshifts $z = 0.5 - 0.8$. The evolution of massive clusters depends exponentially on the amplitude of mass fluctuations and thus provides a powerful measure of this important cosmological parameter. The relatively high abundance of massive clusters observed at $z > 0.5$, and the relatively slow evolution of their abundance with time, suggest a high amplitude of mass fluctuations: $\sigma_8 = 0.9 \pm 10\%$ for $\Omega_m = 0.4$, increasing slightly to $\sigma_8 = 0.95$ for $\Omega_m = 0.25$ and $\sigma_8 = 1.0$ for $\Omega_m = 0.1$ (flat CDM models). We use the cluster abundance observed at $z = 0.5 - 0.8$ to derive a normalization relation from the high-redshift clusters, which is only weakly dependent on Ω_m: $\sigma_8 \Omega_m^{0.14} = 0.78 \pm 0.08$. When combined with recent constraints from the present-day cluster mass function, $\sigma_8 \Omega_m^{0.6} = 0.33 \pm 0.03$, we find $\sigma_8 = 0.98 \pm 0.1$ and $\Omega_m = 0.17 \pm 0.05$. Low σ_8 values ($\lesssim 0.7$) are unlikely; they produce an order of magnitude fewer massive clusters than observed.

Subject headings: cosmological parameters — galaxies: clusters: general

1. Introduction

The amplitude of mass fluctuations is a fundamental cosmological parameter that describes the normalization of the linear spectrum of mass fluctuations in the early universe – the spectrum that seeded galaxy formation. The abundance of massive clusters depends exponentially on this parameter (assuming Gaussian initial fluctuations), because a high amplitude of mass fluctuations forms structure rapidly at early times, while a lower amplitude forms structure more slowly. The most massive systems ($\sim 10^{15} h^{-1} M_\odot$), which take the longest time to form and grow, did not exist at early times if the initial amplitude of mass fluctuations is low, but rather formed only recently.

The amplitude parameter, denoted σ_8 when referring to the *rms* linear density fluctuation in spheres of radius $8h^{-1}$Mpc at $z = 0$, is not easily determined since the mass distribution cannot be directly observed. As a result, this parameter is not yet accurately known. Recent observations suggest an amplitude that ranges in value from $\sigma_8 \sim 0.7$ to a 'high' value of $\sigma_8 \sim 0.9 - 1$. While the difference in the reported values is only around 50%, the impact on structure formation and evolution is much larger, since the latter depends exponentially on σ_8^2. The low amplitude values are suggested by current observations of the CMB spectrum of fluctuations (Netterfield et al. 2002; Sievers et al. 2002; Bond et al. 2002; Ruhl et al. 2002). However, this σ_8 determination is degenerate with the unknown optical depth at reionization: if the optical depth were underestimated, then σ_8 would be as well[1]. Recent observations of the present-day cluster abundance as well as cosmic shear lensing measurements have also suggested that $\sigma_8 \sim 0.7$ (e.g. Jarvis et al. 2003; Hamana et al. 2002; Seljak 2001). However, these measures provide a degenerate relation between the amplitude σ_8 and the mass-density parameter Ω_m: $\sigma_8 \Omega_m^{0.6} \approx 0.33$ (Ikebe et al. 2002; Bahcall et al. 2003; Jarvis et al. 2003; Seljak 2001). The amplitude $\sigma_8 \sim 0.7$ is implied only if $\Omega_m \sim 0.3$. If $\Omega_m \sim 0.2$, as is suggested by some observations (e.g. Carlberg, Yee & Ellingson 1997; Bahcall & Fan 1998; Bahcall et al. 2000; Wilson, et al. 2001; Ikebe et al. 2002; Reiprich & Bohringer 2002), then the amplitude is $\sigma_8 \sim 0.9 - 1$. Early results from the Sloan Digital Sky Survey (SDSS) cluster data (Bahcall et al. 2003) use the shape of the observed cluster mass function to break the degeneracy between the parameters and find $\sigma_8 = 0.9^{+0.3}_{-0.2}$ and $\Omega_m = 0.19^{+0.08}_{-0.07}$. Similar results have recently been obtained from the temperature function of a large sample of X-ray clusters (Ikebe et al. 2002; Reiprich & Bohringer 2002). Most of the recent cluster normalization observations, as well as cosmic shear lensing measurements suggest $\sigma_8 \simeq 0.9 - 1$ if $\Omega_m \simeq 0.2$ (Jarvis et al. 2003; Hamana et al. 2002, and the references above). Combining current CMB measurements with the SDSS cluster mass function yields intermediate values of $\sigma_8 = 0.76 \pm 0.09$ and $\Omega_m = 0.26^{+0.06}_{-0.07}$ (Melchiorri et al. 2003).

The evolution of cluster abundance with time, especially for the most massive clusters, breaks the degeneracy between σ_8 and Ω_m (e.g. Peebles, Daly & Juszkiewicz 1989; Eke, Cole & Frenk 1996; Oukbir & Blanchard 1997; Bahcall, Fan, & Cen 1997; Carlberg et al. 1997; Bahcall & Fan 1998; Donahue & Voit 1999; Henry 2000). This evolution depends strongly on σ_8, and only weakly on Ω_m or other parameters. The expected abundance of massive clusters at $z \sim 0.5 - 1$ differs between Gaussian models with $\sigma_8 = 0.6$ and $\sigma_8 = 1$ by orders-of-magnitude, nearly independently of other parameters (Fan, Bahcall & Cen 1997, e.g.). Therefore, this method provides a uniquely powerful tool in estimating the amplitude σ_8.

[1]See note at end of paper

In this paper we use the abundance of the most massive clusters observed at $z \sim 0.5-0.8$ to place a strong limit on σ_8.

2. The Evolution of Cluster Abundance

A flat $\Omega_m = 0.3$ cold-dark-matter universe (LCDM) with $\sigma_8 = 0.6$ predicts $\sim 10^3$ fewer massive clusters (with mass $\sim 10^{15} h^{-1} M_\odot$) at $z \approx 0.8$ than a universe with $\sigma_8 = 1$ (e.g. Bahcall & Fan 1998; Bode et al. 2001). We use massive clusters observed at $z \approx 0.5$ and $z \approx 0.8$ (as compiled by Bahcall & Fan 1998) to set limits on σ_8. These three clusters, detected originally in X-rays by the EMSS survey (Henry et al. 1992; Luppino & Gioia 1995), have masses larger than $8 \times 10^{14} h^{-1} M_\odot$ within a radius of $1.5 h^{-1}$ comoving Mpc. The masses are determined from gravitational weak lensing observations (for two of the three clusters), as well as from the observed temperatures ($T \gtrsim 8$ Kev) and velocity dispersions ($\gtrsim 1200$ Km/s) of the clusters. All clusters have a measured S-Z effect (Grego 2001; Carlstrom et al. 2001). These clusters have been conservatively selected, with mass measurements available from several independent methods, all yielding consistent results (see Bahcall & Fan (1998) for details, including the consistency of the mass determinations from different methods, and the relevant abundances at $z \approx 0.6$ and 0.8). Since only a threshold cluster mass is used in the analysis below (i.e., not individual cluster masses), a conservative mass threshold of $8 \times 10^{14} h^{-1} M_\odot$ is used; the clusters are well above this threshold. The resulting abundances are $n_{cl} = 1.4^{+1.1}_{-0.9} \times 10^{-8} h^3 \mathrm{Mpc}^{-3}$ at $z = 0.5 - 0.65$, and $1.4^{+1.4}_{-1.1} \times 10^{-8} h^3 \mathrm{Mpc}^{-3}$ at $z = 0.65 - 0.9$. The error bars represent 68% confidence level assuming Poisson statistics and equal likelihood for each $\log(n_{cl})$. The volume searched and hence the number density depends on the assumed cosmology; these numbers are for an LCDM model with $\Omega_m = 0.3$.

For purposes of comparison, the abundance of massive clusters at $z \approx 0$ and at $z \approx 0.38$ is obtained from the temperature function observed by Ikebe et al. (2002) (for $z \approx 0$) and Henry (2000) (for $z \approx 0.38$). Here we convert our threshold mass of $8 \times 10^{14} h^{-1} M_\odot$ (within $1.5 h^{-1}$ comoving Mpc) to temperature using the mean relation between cluster mass (observed directly from weak gravitational lensing) and temperature: $M(\leq 1 h^{-1}$ Mpc$) = 0.95 T(\mathrm{kev})\ 10^{14} h^{-1} M_\odot$ (Bahcall & Sette (2002); see also Hjorth, Oukbir & van Kampen (1998); Carlberg, Yee & Ellingson (1997)). These observations indicate that cluster masses determined from lensing are consistent on average with those derived from X-ray temperatures, with an *rms* scatter of $\sim$20%. For the small extrapolation to radius $1.5 h^{-1}$ Mpc comoving, we use the observed cluster profile on these scales: $M(< R) \sim R^{0.6}$ (Carlberg, Yee & Ellingson 1997; Fischer & Tyson 1997). This allows us to determine the observed cluster abundance at $z \simeq 0$ and $z \simeq 0.38$ for the relevant mass clusters. We find

$n_{cl} = 1.1^{+1.1}_{-0.7} \times 10^{-7} h^3 \mathrm{Mpc}^{-3}$ at $z = 0.05$ (Ikebe et al. 2002) and $n_{cl} = 1.7^{+1.7}_{-1.1} \times 10^{-8} h^3 \mathrm{Mpc}^{-3}$ at $z = 0.3 - 0.5$ (Henry 2000); the error bars allow for statistical uncertainty as well as the uncertainty in the mass threshold. These numbers assume an $\Omega_m = 0.3$ LCDM cosmology.

The observed abundance of these massive clusters as a function of redshift is presented in Figure 1. In the next section, we compare the data with semianalytic predictions and constrain the allowed range of parameters. First, we use only the high redshift ($z > 0.5$) cluster abundance (where mass thresholds are determined by multiple methods, as discussed above). Then we take advantage of the full evolution of the abundance, from $z \sim 0$ to $z \sim 0.8$, by combining the high-redshift result with the independent $z \approx 0$ cluster normalization relation obtained from other studies. We discuss these analyses in the following section.

3. Comparison with Predicted Densities

For a given cosmological model, the expected cluster mass function can be predicted using recent improvements to the Press-Schechter formalism. We follow the procedure outlined in the Appendix of Hu & Kravtsov (2002). For a given choice of parameters, the linear matter power spectrum is calculated with the publicly available CMBFAST code (Seljak & Zaldarriaga 1996). This allows a prediction of the mass function, using a fixed mean overdensity as the definition of mass. This mass can then be extrapolated to 1.5 h^{-1}Mpc comoving, assuming an NFW density profile. The analytic predictions were compared to the N-body results of Bode et al. (2001), and the resulting cluster abundances were found to agree within 20% out to redshift $z = 1$.

The first result is presented in Figure 1, where we compare the observed cluster abundance as a function of redshift with that expected from LCDM models. Here we use LCDM with the 'concordance' value of $\Omega_m = 0.3$ (Bahcall et al. 1999, with $h = 0.72$ and $n = 1$), but with different amplitudes σ_8 ranging from 0.6 to 1.2 (bottom to top curves). Comparison of data and models shows that if $\Omega_m = 0.3$, then σ_8 has to be relatively high, $\sigma_8 = 0.9 \pm 0.1$, in order to produce the observed abundance of clusters at all redshifts. Low normalization values of $\sigma_8 \approx 0.7$— which would be required for $\Omega_m = 0.3$ by the present-day cluster abundance and the cosmic shear lensing measurements discussed in §2— produce an order-of-magnitude too few massive clusters as compared with observations at any redshift. Therefore, for Ω_m=0.3, σ_8 needs to be $\gtrsim 0.9$.

What if Ω_m is not 0.3? How does this affect the allowed range of σ_8? In Figure 2 we compare the data to the cluster evolution expected for $\sigma_8 = 1.1$, 0.9, and 0.7 (top to bottom bands), where the width of each band covers all values of Ω_m from 0.1 to 0.4. It can be seen

that the best fit to the data is again $\sigma_8 \simeq 0.9 - 1$. Low values of $\sigma_8 \lesssim 0.7$ appear to be excluded by the data, nearly independent of Ω_m; they provide significantly fewer high mass clusters than observed, especially at $z > 0.5$. This result is consistent with the independent SDSS cluster mass-function at low redshift which yields $\sigma_8 = 0.9^{+0.3}_{-0.2}$ and $\Omega_m = 0.19^{+0.08}_{-0.07}$ (Bahcall et al. 2003). The independent result obtained here from the high redshift clusters (Figures 1-2) provides important confirmation that σ_8 is indeed high ($\sim 0.9 - 1$).

Finally, we use the observed cluster abundance at high redshifts ($z \gtrsim 0.5$), independent of the lower redshift points, to determine the best $\sigma_8 - \Omega_m$ normalization relation from high redshift clusters. This was done by minimizing χ^2, using the error bars given in Bahcall & Fan (1998) plus an additional factor of 20% to account for uncertainties in the analytic prediction. The allowed 68% and 95% confidence limits are presented by the solid contours in Figure 3. As expected, the observed abundance of these massive clusters at high redshift depends mostly on σ_8 (§2); the dependence on Ω_m is very weak. (The dependence on other parameters - Hubble constant and spectral index – is also weak, as is discussed below). This fact makes the high redshift cluster abundance method a powerful tool in constraining σ_8. The results in Figure 3 show that for any observationally acceptable range of Ω_m, from ~ 0.1 to ~ 0.4, the amplitude remains in the range of $\sigma_8 \simeq 0.9 - 1$. Values of $\sigma_8 < 0.8$ are unlikely; they have too little power to form massive systems at $z > 0.5$.

We use the $z = 0.5 - 0.8$ clusters to determine the best-fit cluster normalization relation from the high-redshift objects; we find:

$$\sigma_8 = 1.03 - 0.3\Omega_m \pm 10\% \qquad \text{, or} \tag{1a}$$

$$\sigma_8 \Omega_m^{0.14} = 0.78 \pm 0.08 \tag{1b}$$

(where σ_8 and Ω_m refer, as before, to their $z = 0$ value). The linear relation [1a] is within 3% of the true best-fit σ_8 over the range $\Omega_m = 0.1 - 1.0$; the power-law fit [1b] is superior to equation [1a] in the range $\Omega_m = 0.2 - 0.7$. These relations assume a Hubble constant of $h = 0.72$ and spectral index $n = 1$, but the results are insensitive to reasonable changes in h and n. We find that changing h by $\Delta h = \pm 0.13$, i.e., from $h = 0.59$ to $h = 0.85$, changes σ_8 by less than $\pm 5\%$ (at a given Ω_m); σ_8 increases slightly with h. Similarly, changing the spectral index $n = 1$ by $\Delta n = \pm 0.2$, from $n = 0.8$ to $n = 1.2$, changes σ_8 by less than $\pm 5\%$ (in the same direction: slightly higher σ_8 for higher n). The 1- and 2-sigma contours of the allowed σ_8 - Ω_m parameter region that includes both these h and n variations are presented by the dotted curves in Figure 3. As expected, these conservative ranges in h and n broaden the allowed region. The above results include the estimated uncertainty in the mass threshold for the clusters (Bahcall & Fan 1998). As discussed in §2, only a lower threshold (selected at the lower 1-sigma level of the mass estimates) is used in this analysis, not the individual mass of each cluster. Reducing the cluster mass threshold further by 10%

will reduce the amplitude σ_8 by $\sim$10% (Fan, Bahcall & Cen 1997; Bahcall & Fan 1998).

Also shown in Figure 3 are the analogous confidence contours from the cluster mass-function normalization at $z = 0.1 - 0.2$ (Bahcall et al. 2003); the best-fit relation is approximated by

$$\sigma_8 \Omega_m^{0.6} = 0.33 \pm 0.03 \tag{2}$$

which has a steeper Ω_m dependence than the high-redshift constraint. These two independent constraints overlap only at low Ω_m. Requiring that both of the constraints of equations [1] and [2] be simultaneously satisfied yields

$$\sigma_8 = 0.98 \pm 0.1 \tag{3a}$$

$$\Omega_m = 0.17 \pm 0.05 \tag{3b}$$

for the allowed 1-sigma overlap region when h=0.72 and n=1. Allowing for variations in h and n as discussed above yields the same central values for σ_8 and Ω_m, but broadens the allowed range. Even with the very broad ranges adopted for h and n, the conservative limits $\sigma_8 > 0.70$ and $\Omega_m < 0.36$ can be set at the 95% confidence level.

4. Discussion and Conclusions

We use the observed abundance of high-mass clusters of galaxies at $z = 0.5 - 0.8$ to determine the linear amplitude of mass fluctuations, σ_8. The cluster abundance depends exponentially on this amplitude, and only weakly on other parameters; it therefore provides a powerful method for measuring this important parameter. We show that the relatively high abundance of massive clusters observed at $z \gtrsim 0.5$, as well as their relatively slow evolution with time, requires a high amplitude of mass fluctuations, $\sigma_8 \sim 0.9 - 1$. This conclusion is nearly independent of the exact value of Ω_m (in the typical range of $\Omega_m \sim 0.1 - 0.4$).

We use the observed abundance at $z \gtrsim 0.5$ to determine a normalization relation from high redshift clusters. The relation depends only weakly on Ω_m: $\sigma_8 \Omega_m^{0.14} = 0.78 \pm 0.08$; alternatively, a linear relation of the form $\sigma_8 = 1.03 - 0.3\Omega_m$ ($\pm$10%) provides a similarly good fit to the data. These fits illustrate that $\sigma_8 \gtrsim 0.8$ for any $\Omega_m \leq 0.4$. For the typical observationally suggested value of $\Omega_m \simeq 0.2 - 0.3$, the amplitude is $\sigma_8 = 0.95 \pm 0.1$. We emphasize that this high σ_8 value indicated by the cluster abundance at high redshift is nearly independent of the exact value of Ω_m.

We combine the high redshift constraint above with the independent normalization relation obtained from low redshift cluster abundance— a relation that is steeper in Ω_m

($\sim \Omega_m^{0.6}$; equation [2]). The combination breaks the degeneracy between the two parameters. We find $\sigma_8 = 0.98 \pm 0.1$ and $\Omega_m = 0.17 \pm 0.05$ (Figure 3; equation [3]).

The high value of σ_8 required to explain the high abundance of the most massive clusters at $z \gtrsim 0.5$ is consistent with the present day cluster mass function if the mass density parameter is low, $\Omega_m \sim 0.2$. If Ω_m=0.3, then the high redshift clusters still require a high amplitude ($\sigma_8 = 0.92 \pm 0.1$), since this constraint is nearly independent of Ω_m; the low redshift cluster abundance is consistent with this value at the 2-sigma level.

These results improve upon Bahcall & Fan (1998) by using improvements over the standard Press-Schechter formula (which does not accurately reproduce results from cosmological simulations at high redshift), allowing for changes in h and n, and by using a more recent cluster normalization relation at low redshift. The current results yield slightly lower values for the cosmological parameters than the previous work (the latter suggested $\sigma_8 = 1.2 \pm 0.22$ and $\Omega_m = 0.2^{+0.13}_{-0.07}$ (68%) when using the most massive clusters) but are consistent with the new values within the error-bars (Bahcall & Fan 1998; Fan, Bahcall & Cen 1997).

The excess CMB fluctuations detected on small scales by the CBI (Mason et al. 2002) and the BIMA (Dawson et al. 2002) experiments implies (if correctly interpreted as being due to the S-Z effect from distant clusters) that $\sigma_8 = 1.04 \pm 0.12$ (95%; Komatsu & Seljak 2002). This is in excellent agreement with our current conclusions. We note, however, that this high amplitude is inconsistent with the lower value of $\sigma_8 \approx 0.7$ suggested by current CMB data on large scales (which is degenerate with the unknown optical depth). Future CMB observations should clarify this current inconsistency. If massive clusters exist with relatively high abundance at high redshifts, as suggested by the data used here (as well as by deep X-ray surveys, e.g. Rosati, Borgani & Norman 2002), then these clusters should indeed produce the excess S-Z fluctuations observed by the CMB data.

The relatively high abundance of massive clusters observed at $z \gtrsim 0.5$ provides one of the strongest arguments for a high amplitude of mass fluctuations, $\sigma_8 \simeq 1$.

Note added March 7, 2003: The recent CMB anisotropy spectrum released by the WMAP team in February 2003 nicely confirms the results presented here. The constraint from the CMB alone is $\sigma_8 = 0.9 \pm 0.1$ (Spergel et al. 2003), in full agreement with the current high redshift cluster constraint.

This research was supported by the National Computational Science Alliance under NSF Cooperative Agreement ASC97-40300, PACI Subaward 766.

REFERENCES

Bahcall, N.A. & Fan, X. 1998, ApJ, 504, 1

Bahcall, N.A., Fan, X. & Cen, R. 1997, ApJ, 485, L53

Bahcall, N.A. & Sette, A. 2002, Junior Thesis, Princeton University

Bahcall, N. A., Ostriker, J. P., Perlmutter, S. & Steinhardt, P. 1999, Science, 284, 1481

Bahcall, N. A., Cen, R., Davé, R., Ostriker, J. P. & Yu, Q. 2000, ApJ, 541, 1

Bahcall, N.A., et al. 2003, ApJ, in press (astro-ph/0205490)

Bode, P., Bahcall, N.A., Ford, E.B., Ostriker, J.P. 2001, ApJ, 551, 15

Bond, J.R., et al. 2002, ApJ, submitted (astro-ph/0205386)

Carlberg, R.G., Morris, S.L., Yee, H.K.C. & Ellingson, E. 1997, ApJ, 479, L19

Carlberg, R.G., Yee, H.K.C. & Ellingson, E. 1997, ApJ, 478, 462

Carlstrom, J.E. et al. 2001, in Constructing the Universe with Clusters of Galaxies, IAP Conference Proceedings, ed. F. Durret & G. Gerbal, preprint (astro-ph/0103480)

Dawson, K.S., et al. 2002, ApJ, in press (astro-ph/0206012)

Donahue, M. & Voit, G.M. 1999, ApJ, 523, L137

Eke, V.R., Cole, S. & Frenk C.S. 1996, MNRAS, 282, 263

Fan, X., Bahcall, N.A. & Cen, R. 1997, ApJ, 490, L123

Fischer, P. & Tyson, J. A. 1997, AJ, 1 14, 14

Grego, L. 2001 ApJ, 552, 2

Hamana, T., et al. 2002, ApJ, submitted (astro-ph/0210450)

Henry, J.P., Gioia, I.M., Maccacaro, T., Morris, S.L., Stocke, J.T. & Wolter, A. 1992 ApJ, 386, 408

Henry, J.P. 2000 ApJ, 534, 565

Hjorth, J., Oukbir, J. & van Kampen, E. 1998, MNRAS, 298, L1

Hu, W. & Kravtsov, A.V. 2002, ApJ, submitted (astro-ph/0203169)

Ikebe, Y., Reiprich, T. H., B?hringer, H., Tanaka, Y. & Kitayama, T. 2002, A&A, 383, 773

Jarvis, M., et al. 2003, AJ, in press (astro-ph/0210604)

Komatsu, E. & Seljak, U. 2002, MNRAS, 336, 1256

Luppino, G.A. & Gioia, I.M. 1995, ApJ, 445, L77

Mason, B.S., et al. 2002, ApJ, submitted (astro-ph/0205384)

Melchiorri, A., Bode, P., Bahcall, N.A., & Silk, J. 2003, ApJ, in press (astro-ph/0212276)

Netterfield, C.B., et al. 2002, ApJ, 571, 604

Oukbir, J. & Blanchard, A. 1997, A&A, 317, 1

Peebles, P.J.E., Daly, R.A. & Juszkiewicz, R. 1989, ApJ, 347, 563

Reiprich, T. H. & Bohringer, H. 2002, ApJ, 567, 716

Rosati, P., Borgani, S. & Norman, C. 2002, ARA&A, 40, 539

Ruhl, J.E., et al. 2002, ApJ, submitted (astro-ph/0212229)

Seljak, U., MNRAS, submitted (astro-ph/0111362)

Seljak, U. & Zaldarriaga, M. 1996, ApJ, 469, 437

Sievers, J.L., et al. 2002, ApJ, submitted (astro-ph/0205387)

Spergel, D.N., et al. 2003, ApJ, submitted (astro-ph/0302209)

Wilson, G., Kaiser, N., Luppino, G. & Cowie, L.L. 2001, ApJ, 555, 572

This preprint was prepared with the AAS LaTeX macros v5.0.



Fig. 1.— Evolution of cluster abundance with redshift, for clusters with mass $M_{1.5,com} \geq 8 \times 10^{14} h^{-1} M_\odot$ (within a comoving radius of $1.5 h^{-1}$Mpc). Dots with error bars are the data as described in the text. The lines are the predicted number density assuming a flat $\Omega_m = 0.3$ cosmology; each line is labeled with the σ_8 used.



Fig. 2.— Same as Figure 1, but for $0.1 \leq \Omega_m \leq 0.4$. Each band is labeled with the σ_8 used to predict the number density, and within each band Ω_m is varied from 0.1 (bottom of band) to 0.4 (top); $h = 0.72$ and $n = 1$.



Fig. 3.— Confidence contours in the $\Omega_m - \sigma_8$ plane. Solid lines are the 68% and 95% confidence contours found from the high-redshift cluster abundance (contours extending to high Ω_m) and from the low-redshift SDSS HMF clusters (extending to lower σ_8); these assume $h = 0.72$ and $n = 1$ and two degrees of freedom. The shaded region satisfies both constraints at the 68% level. The dotted lines are the 68% and 95% limits with four degrees of freedom when allowing $0.59 < h < 0.85$ and $0.8 < n < 1.2$.